# Management of Residents with Diarrhea in Continuing Care

Central Zone Infection Prevention and Control (IPC)



#### Suspect Gastrointestinal Illness (GI)

2 or more episodes of fluffy pieces in liquid or watery stool in a 24-hour period, which is new or unusual for the resident

OR

1 episode of bloody diarrhea

#### Initiate Precautions



- ✓ Collect stool sample and document frequency and consistency of bowel movements
- ✓ Refer to the Continuing Care Resource Manual

## Does the patient have loose or watery stool?

Fluffy pieces in liquid or watery stool (Type 6 or 7 on the Bristol Stool Chart)

Do you need to consider a GI/CDI outbreak?

2 or more people with onset of similar symptoms in a 48-hour period

Refer to MOH - Central Zone | Alberta Health Services for additional

resources

### Suspect Clostridioides difficile Infection (CDI)

**3 or more episodes** of fluffy pieces in liquid or watery stool in a **24-hour** period, which is new or unusual for the resident

OR

**6 or more** fluffy pieces in liquid or watery stool in a **36-hour** period, which is new or unusual for the resident

Initiate Precaution

CONTACT PRECAUTIONS
SPORICIDAL CLEAN

STOP

- Collect stool sample and document frequency and consistency of bowel movements
- ✓ Refer to CDI Management on Central Zone tab for further information
- ✓ Refer to the Continuing Care Resource Manual

Discontinue precautions when resident has been without symptoms for 48 hours and has had at least one normal or formed bowel movement

For more information contact: <u>ipccenzadmin@ahs.ca</u>

© 2023 Alberta Health Services

(cc) BY-NC-SA

https://creativecommons.org/licenses/by-nc-sa/4.0/

Original date: October 2016 Revised date: November 2023

